arXiv:1402.6080v1 [math.FA] 25 Feb 2014

COMPARISON OF THE SPEED OF CONVERGENCE AMONG VARIOUS ITERATIVE SCHEMES

VATAN KARAKAYA, FAIK GÜRSOY, AND MÜZEYYEN ERTÜRK

Abstract. We show that iterative scheme due to Karahan and Özdemir (2013) can be used to approximate fixed point of contraction mappings. Furthermore, we prove that CR iterative scheme converges faster than the iterative scheme due to Karahan and Özdemir (2013) for the class contraction mappings. Finally, we prove a data dependence result for contraction mappings by employing iterative scheme due to Karahan and Özdemir (2013).

1. introduction

Fixed point theory has been appeared as one of the most powerful and substantial theoretical tools of mathematics. This theory has a long history and has been studied intensively by many researchers in various aspects. For the past 30 years or so, the study of iterative procedures for the approximation of fixed points of various classes of operators have been flourishing areas of research for many mathematicians. Consequently, considerable research efforts have been devoted to introduce various iteration methods and study its more qualitative features, for example, [1, 8, 12, 15, 16, 17, 18, 19, 20, 22, 23, 26, 29, 32].

We begin our exposition with an overview of various iterative methods.

Throughout this paper $\mathbb{N}$ denotes set of all nonnegative integers including zero. Let B be a Banach space, S be a subset of B and T be a selfmap of S. Let $\left\{\alpha_n^i\right\}_{n=0}^{\infty}$, $i \in \{1,2,3\}$ be real sequences in $[0,1]$ satisfying certain control condition(s).

An iterative sequence $\{x_n\}_{n=0}^{\infty}$ defined by

$$\begin{cases} x_0 \in S, \\ x_{n+1} = Tx_n,\ n \in \mathbb{N}, \end{cases} \tag{1.1}$$

is known as Picard iteration procedure [23], which is commonly used to approximate fixed point of contraction mappings satisfying

$$\|Tx - Ty\| \leq \delta \|x - y\|,\ \delta \in (0,1),\ \text{for all } x, y \in B. \tag{1.2}$$

The following iteration methods are called Noor [18], and SP [22] iteration methods, respectively:

$$\begin{cases} x_0 \in S, \\ x_{n+1} = \left(1-\alpha_n^1\right) x_n + \alpha_n^1 T y_n, \\ y_n = \left(1-\alpha_n^2\right) x_n + \alpha_n^2 T z_n, \\ z_n = \left(1-\alpha_n^3\right) x_n + \alpha_n^3 T x_n,\ n \in \mathbb{N}, \end{cases} \tag{1.3}$$

2000 *Mathematics Subject Classification.* Primary 47H06, 54H25.

Key words and phrases. Iterative schemes, Convergence, Speed of convergence, Equivalence of convergence, Data dependence of fixed points.

and

$$\begin{cases} x_0 \in S, \\ x_{n+1} = \left(1-\alpha_n^1\right) y_n + \alpha_n^1 T y_n, \\ y_n = \left(1-\alpha_n^2\right) z_n + \alpha_n^2 T z_n, \\ z_n = \left(1-\alpha_n^3\right) x_n + \alpha_n^3 T x_n,\ n \in \mathbb{N}. \end{cases} \tag{1.4}$$

Remark 1. *(i) Noor iteration method (1.5) reduces to well-known Ishikawa iteration method* [12] *for* $\alpha_n^3 = 0$ *and Mann iteration method* [17] *for* $\alpha_n^2 = \alpha_n^3 = 0$. *(ii) SP iteration method (1.6) reduces to a two-step Mann iteration method* [32] *for* $\alpha_n^3 = 0$.

Agarwal et al. [1] inroduced an S-iteration method as follows

$$\begin{cases} s_0 \in S, \\ s_{n+1} = \left(1-\alpha_n^1\right) T s_n + \alpha_n^1 T t_n, \\ t_n = \left(1-\alpha_n^2\right) s_n + \alpha_n^2 T s_n,\ n \in \mathbb{N}. \end{cases} \tag{1.5}$$

The following iteration method is referred to as CR iteration method [9]

$$\begin{cases} u_0 \in S, \\ u_{n+1} = \left(1-\alpha_n^1\right) v_n + \alpha_n^1 T v_n, \\ v_n = \left(1-\alpha_n^2\right) T u_n + \alpha_n^2 T y_n, \\ y_n = \left(1-\alpha_n^3\right) u_n + \alpha_n^3 T u_n,\ n \in \mathbb{N}. \end{cases} \tag{1.6}$$

Very recently, Karahan and Özdemir [13] introduced a new three step iteration as follows

$$\begin{cases} p_0 \in S, \\ p_{n+1} = \left(1-\alpha_n^1\right) T p_n + \alpha_n^1 T q_n, \\ q_n = \left(1-\alpha_n^2\right) T p_n + \alpha_n^2 T r_n, \\ r_n = \left(1-\alpha_n^3\right) p_n + \alpha_n^3 T p_n,\ n \in \mathbb{N}. \end{cases} \tag{1.7}$$

Convergence analysis of iterative methods has an important role in the study of iterative approximation of fixed point theory. Fixed point iteration methods may exhibit radically different behaviors for various classes of mappings. While a particular fixed point iteration method is convergent for an appropriate class of mappings, it may not be convergent for the others. Due to various reasons, it is important to determine whether an iteration method converges to fixed point of a mapping. In many cases, there can be two or more than two iteration procedures approximating to a fixed point of a mapping, for example, [7, 14, 25, 31]. In such cases, the critical and important point is to compare rate of convergence of these iterations to find out which ones converge faster to that fixed point, e.g., [2, 4, 5, 6, 10, 11, 21, 24, 27, 28, 34, 35].

Recently, several authors introduced different type iteration methods and they have proved that their iteration methods converges faster than Picard, Mann and Ishikawa iteration methods, e.g., [9, 13, 14, 21, 28].

In this paper, we are concerned with two recent iteration methods defined by (1.6) and (1.7). We show that iteration method (1.7) converges to fixed point of a contraction mapping satisfying (1.2). Also, we prove that CR iteration method (1.6) is equivalent and faster than iteration method (1.7) for the class of contraction mappings. Finally, we give a data dependence result for the fixed point of contraction mappings using iteration method (1.7).

In order to obtain our main results we need following lemmas and definitions.

Definition 1. [3] *Let $\{a_n\}_{n=0}^{\infty}$ and $\{b_n\}_{n=0}^{\infty}$ be two sequences of real numbers with limits a and b, respectively. Assume that there exists*

$$\lim_{n\to\infty}\frac{|a_n-a|}{|b_n-b|}=l. \tag{1.8}$$

(i) If $l=0$, the we say that $\{a_n\}_{n=0}^{\infty}$ converges faster to a than $\{b_n\}_{n=0}^{\infty}$ to b.

(ii) If $0<l<\infty$, then we say that $\{a_n\}_{n=0}^{\infty}$ and $\{b_n\}_{n=0}^{\infty}$ have the same rate of convergence.

Definition 2. [3] *Let T,$\widetilde{T}:B\to B$ be two operators. We say that $\widetilde{T}$ is an approximate operator of T if for all $x\in B$ and for a fixed $\varepsilon>0$ we have*

$$\left\|Tx-\widetilde{T}x\right\|\leq\varepsilon. \tag{1.9}$$

Lemma 1. [33]*Let $\{a_n\}_{n=0}^{\infty}$ and $\{\rho_n\}_{n=0}^{\infty}$ be nonnegative real sequences satisfying the following inequality:*

$$a_{n+1}\leq(1-\eta_n)\,a_n+\rho_n, \tag{1.10}$$

where $\eta_n\in(0,1)$, for all $n\geq n_0$, $\sum_{n=1}^{\infty}\eta_n=\infty$, and $\frac{\rho_n}{\eta_n}\to 0$ as $n\to\infty$. Then $\lim_{n\to\infty}a_n=0$.

Lemma 2. [30] *Let $\{a_n\}_{n=0}^{\infty}$ be a nonnegative sequence for which one assumes there exists $n_0\in\mathbb{N}$, such that for all $n\geq n_0$ one has satisfied the inequality*

$$a_{n+1}\leq(1-\mu_n)\,a_n+\mu_n\eta_n, \tag{1.11}$$

where $\mu_n\in(0,1)$, for all $n\in\mathbb{N}$, $\sum\limits_{n=0}^{\infty}\mu_n=\infty$ and $\eta_n\geq 0$, $\forall n\in\mathbb{N}$. Then the following inequality holds

$$0\leq\limsup_{n\to\infty}a_n\leq\limsup_{n\to\infty}\eta_n. \tag{1.12}$$

2. Main Results

Theorem 1. *Let S be a nonempty closed convex subset of a Banach space B and $T:S\to S$ be a contraction map satisfying condition (1.2). Let $\{p_n\}_{n=0}^{\infty}$ be an iterative sequence generated by (1.7) with real sequences $\left\{\alpha_n^i\right\}_{n=0}^{\infty}$, $i\in\{1,2,3\}$ in $[0,1]$ satisfying $\sum_{k=0}^{n}\alpha_k^1=\infty$. Then $\{p_n\}_{n=0}^{\infty}$ converges to a unique fixed point of T, say x_*.*

Proof. Picard-Banach theorem guarantees the existence and uniqueness of x_*. We will show that $p_n\to x_*$ as $n\to\infty$. From (1.2) and (1.7) we have

$$\begin{aligned}\|r_n-x_*\| &= \left\|\left(1-\alpha_n^3\right)p_n+\alpha_n^3Tp_n-\left(1-\alpha_n^3+\alpha_n^3\right)x_*\right\|\\ &\leq \left(1-\alpha_n^3\right)\|p_n-x_*\|+\alpha_n^3\|Tp_n-Tx_*\|\\ &\leq \left(1-\alpha_n^3\right)\|p_n-x_*\|+\alpha_n^3\delta\|p_n-x_*\|\\ &= \left[1-\alpha_n^3(1-\delta)\right]\|p_n-x_*\|, \qquad (2.1)\end{aligned}$$

$$\begin{aligned}\|q_n-x_*\| &\leq \left(1-\alpha_n^2\right)\|Tp_n-Tx_*\|+\alpha_n^2\|Tr_n-Tx_*\|\\ &\leq \left(1-\alpha_n^2\right)\delta\|p_n-x_*\|+\alpha_n^2\delta\|r_n-x_*\|\\ &\leq \left\{\left(1-\alpha_n^2\right)\delta+\alpha_n^2\delta\left[1-\alpha_n^3(1-\delta)\right]\right\}\|p_n-x_*\|, \qquad (2.2)\end{aligned}$$

and

$$\begin{aligned}\|p_{n+1}-x_*\| &\leq \left(1-\alpha_n^1\right)\|Tp_n-Tx_*\|+\alpha_n^1\|Tq_n-Tx_*\| \\ &\leq \left(1-\alpha_n^1\right)\delta\|p_n-x_*\|+\alpha_n^1\delta\|q_n-x_*\| \\ &\leq \left\{\left(1-\alpha_n^1\right)\delta \right. \\ &\quad \left. +\alpha_n^1\delta\left\{\left(1-\alpha_n^2\right)\delta+\alpha_n^2\delta\left[1-\alpha_n^3\left(1-\delta\right)\right]\right\}\right\}\|p_n-x_*\| \end{aligned} \tag{2.3}$$

Since $\delta\in(0,1)$ and $\alpha_n^i\in[0,1]$, for all $n\in\mathbb{N}$ and for each $i\in\{1,2,3\}$

$$1-\alpha_n^2\left(1-\delta\right)<1, \tag{2.4}$$

$$1-\alpha_n^3\left(1-\delta\right)<1. \tag{2.5}$$

By using $\delta\in(0,1)$, (2.4) and (2.5) in (2.3), we obtain

$$\begin{aligned}\|p_{n+1}-x_*\| &\leq \left[\left(1-\alpha_n^1\right)\delta\left\{\left(1-\alpha_n^2\right)\delta+\alpha_n^2\delta\right\}+\alpha_n^1\delta\right]\|p_n-x_*\| \\ &\leq \left[\left(1-\alpha_n^1\right)\delta\left\{1-\alpha_n^2\left(1-\delta\right)\right\}+\alpha_n^1\delta\right]\|p_n-x_*\| \\ &\leq \left[1-\alpha_n^1\left(1-\delta\right)\right]\|p_n-x_*\| \\ &\leq \cdots \\ &\leq \prod_{k=0}^{n}\left[1-\alpha_k^1\left(1-\delta\right)\right]\|p_0-x_*\|. \end{aligned} \tag{2.6}$$

It is well-known from the classical analysis that $1-x\leq e^{-x}$ for all $x\in[0,1]$. By considering this fact together with (2.6), we obtain

$$\begin{aligned}\|p_{n+1}-x_*\| &\leq \prod_{k=0}^{n}\left[1-\alpha_k^1\left(1-\delta\right)\right]\|p_0-x_*\| \\ &\leq \frac{\|p_0-x_*\|}{e^{(1-\delta)\sum_{k=0}^{n}\alpha_k^1}}. \end{aligned} \tag{2.7}$$

Taking the limit of both sides of inequality (2.7) yields $p_n\to x_*$ as $n\to\infty$. □

Theorem 2. *Let S, B and T with fixed point x_* be as in Theorem 1. Let $\{u_n\}_{n=0}^{\infty}$, $\{p_n\}_{n=0}^{\infty}$be two iterative sequences defined by (1.6) and (1.7) with real sequences $\left\{\alpha_n^i\right\}_{n=0}^{\infty}$, $i\in\{1,2,3\}$ in $[0,1]$ satisfying $\sum_{k=0}^{n}\alpha_k^1=\infty$. Then the following are equivalent:*

(i) the iteration method (1.7) converges to the fixed point x_* of T;

(ii) the CR iteration method (1.6) converges to the fixed point x_* of T.

Proof. We will prove (i)⇒(ii), that is, if iteration method (1.7) converges to x_*, then CR iteration method (1.6) does too. Now by using iteration method (1.7), CR

iteration method (1.6) and condition (1.2), we have

$$\begin{aligned}
\|p_{n+1}-u_{n+1}\| &= \left\|\left(1-\alpha_n^1\right)Tp_n+\alpha_n^1Tq_n-\left(1-\alpha_n^1\right)v_n-\alpha_n^1Tv_n\right\| \\
&\leq \left(1-\alpha_n^1\right)\|Tp_n-v_n\|+\alpha_n^1\|Tq_n-Tv_n\| \\
&\leq \left(1-\alpha_n^1\right)\|p_n-v_n\|+\alpha_n^1\delta\|q_n-v_n\|+\left(1-\alpha_n^1\right)\|p_n-Tp_n\| \\
&= \left(1-\alpha_n^1\right)\left\|\left(1-\alpha_n^2+\alpha_n^2\right)p_n-\left(1-\alpha_n^2\right)Tu_n-\alpha_n^2Ty_n\right\| \\
&\quad +\alpha_n^1\delta\|q_n-v_n\|+\left(1-\alpha_n^1\right)\|p_n-Tp_n\| \\
&\leq \left(1-\alpha_n^1\right)\left(1-\alpha_n^2\right)\|p_n-Tu_n\|+\left(1-\alpha_n^1\right)\alpha_n^2\|p_n-Ty_n\| \\
&\quad +\alpha_n^1\delta\|q_n-v_n\|+\left(1-\alpha_n^1\right)\|p_n-Tp_n\| \\
&\leq \left(1-\alpha_n^1\right)\left(1-\alpha_n^2\right)\{\|p_n-Tp_n\|+\|Tp_n-Tu_n\|\} \\
&\quad +\left(1-\alpha_n^1\right)\alpha_n^2\{\|p_n-Tp_n\|+\|Tp_n-Ty_n\|\} \\
&\quad +\alpha_n^1\delta\|q_n-v_n\|+\left(1-\alpha_n^1\right)\|p_n-Tp_n\| \\
&\leq \left(1-\alpha_n^1\right)\left(1-\alpha_n^2\right)\delta\|p_n-u_n\|+\left(1-\alpha_n^1\right)\left(1-\alpha_n^2\right)\|p_n-Tp_n\| \\
&\quad +\left(1-\alpha_n^1\right)\alpha_n^2\|p_n-Tp_n\|+\left(1-\alpha_n^1\right)\alpha_n^2\delta\|p_n-y_n\| \\
&\quad +\alpha_n^1\delta\|q_n-v_n\|+\left(1-\alpha_n^1\right)\|p_n-Tp_n\| \\
&= \left(1-\alpha_n^1\right)\left(1-\alpha_n^2\right)\delta\|p_n-u_n\| \\
&\quad +\left(1-\alpha_n^1\right)\alpha_n^2\delta\left\|\left(1-\alpha_n^3+\alpha_n^3\right)p_n-\left(1-\alpha_n^3\right)u_n-\alpha_n^3Tu_n\right\| \\
&\quad +\alpha_n^1\delta\|q_n-v_n\|+2\left(1-\alpha_n^1\right)\|p_n-Tp_n\| \\
&\leq \left(1-\alpha_n^1\right)\left(1-\alpha_n^2\right)\delta\|p_n-u_n\| \\
&\quad +\left(1-\alpha_n^1\right)\alpha_n^2\delta\left(1-\alpha_n^3\right)\|p_n-u_n\|+\left(1-\alpha_n^1\right)\alpha_n^2\delta\alpha_n^3\|p_n-Tu_n\| \\
&\quad +\alpha_n^1\delta\|q_n-v_n\|+2\left(1-\alpha_n^1\right)\|p_n-Tp_n\| \\
&\leq \left(1-\alpha_n^1\right)\left(1-\alpha_n^2\right)\delta\|p_n-u_n\|+\left(1-\alpha_n^1\right)\alpha_n^2\delta\left(1-\alpha_n^3\right)\|p_n-u_n\| \\
&\quad +\left(1-\alpha_n^1\right)\alpha_n^2\delta\alpha_n^3\|p_n-Tp_n\|+\left(1-\alpha_n^1\right)\alpha_n^2\delta\alpha_n^3\delta\|p_n-u_n\| \\
&\quad +\alpha_n^1\delta\|q_n-v_n\|+2\left(1-\alpha_n^1\right)\|p_n-Tp_n\| \\
&= \left\{\left(1-\alpha_n^1\right)\left(1-\alpha_n^2\right)\delta+\left(1-\alpha_n^1\right)\alpha_n^2\delta\left[1-\alpha_n^3\left(1-\delta\right)\right]\right\}\|p_n-u_n\| \\
&\quad +\alpha_n^1\delta\|q_n-v_n\|+\left(1-\alpha_n^1\right)\left(2+\alpha_n^2\delta\alpha_n^3\right)\|p_n-Tp_n\|, \qquad (2.8)
\end{aligned}$$

$$\begin{aligned}
\|q_n-v_n\| &= \left\|\left(1-\alpha_n^2\right)Tp_n+\alpha_n^2Tr_n-\left(1-\alpha_n^2\right)Tu_n-\alpha_n^2Ty_n\right\| \\
&\leq \left(1-\alpha_n^2\right)\delta\|p_n-u_n\|+\alpha_n^2\delta\|r_n-y_n\| \\
&= \left(1-\alpha_n^2\right)\delta\|p_n-u_n\| \\
&\quad +\alpha_n^2\delta\left\|\left(1-\alpha_n^3\right)p_n+\alpha_n^3Tp_n-\left(1-\alpha_n^3\right)u_n-\alpha_n^3Tu_n\right\| \\
&\leq \left(1-\alpha_n^2\right)\delta\|p_n-u_n\|+\alpha_n^2\delta\left[1-\alpha_n^3\left(1-\delta\right)\right]\|p_n-u_n\| \\
&\leq \left(1-\alpha_n^2\right)\|p_n-u_n\|+\alpha_n^2\delta\|p_n-u_n\| \\
&= \left[1-\alpha_n^2\left(1-\delta\right)\right]\|p_n-u_n\|. \qquad (2.9)
\end{aligned}$$

Substituting (2.9) in (2.8)

$$\begin{aligned}
\|p_{n+1}-u_{n+1}\| &\leq \left\{\left(1-\alpha_n^1\right)\left(1-\alpha_n^2\right)\delta+\left(1-\alpha_n^1\right)\alpha_n^2\delta\left[1-\alpha_n^3\left(1-\delta\right)\right]\right. \\
&\quad \left.+\alpha_n^1\delta\left[1-\alpha_n^2\left(1-\delta\right)\right]\right\}\|p_n-u_n\| \\
&\quad +\left(1-\alpha_n^1\right)\left(2+\alpha_n^2\delta\alpha_n^3\right)\|p_n-Tp_n\|. \qquad (2.10)
\end{aligned}$$

Since $\delta \in [0,1)$, $\alpha_n^i \in [0,1]$ for all $n \in \mathbb{N}$ and for each $i \in \{1,2,3\}$,

$$1-\alpha_n^2(1-\delta)<1, \tag{2.11}$$

$$1-\alpha_n^3(1-\delta)<1. \tag{2.12}$$

By applying inequalities (2.11) and (2.12) to (2.10), we obtain

$$\begin{aligned} \|p_{n+1}-u_{n+1}\| &\leq \left[1-\alpha_n^1(1-\delta)\right]\|p_n-u_n\| \\ &\quad +\left(1-\alpha_n^1\right)\left(2+\alpha_n^2\delta\alpha_n^3\right)\|p_n-Tp_n\|. \end{aligned} \tag{2.13}$$

Using the fact $x_* = Tx_*$ and triangle inequality for norms, we derive

$$\begin{aligned} \|p_n-Tp_n\| &= \|p_n-x_*+Tx_*-Tp_n\| \\ &\leq \|p_n-x_*\|+\|Tx_*-Tp_n\| \\ &\leq (1+\delta)\|p_n-x_*\|. \end{aligned} \tag{2.14}$$

Substituting (2.14) in (2.13)

$$\begin{aligned} \|p_{n+1}-u_{n+1}\| &\leq \left[1-\alpha_n^1(1-\delta)\right]\|p_n-u_n\| \\ &\quad +\left(1-\alpha_n^1\right)\left(2+\alpha_n^2\delta\alpha_n^3\right)(1+\delta)\|p_n-x_*\|. \end{aligned} \tag{2.15}$$

Denote that

$$\begin{aligned} a_n &= \|p_n-u_n\|, \\ \eta_n &= \alpha_n^1(1-\delta)\in(0,1), \\ \rho_n &= \left(1-\alpha_n^1\right)\left(2+\alpha_n^2\delta\alpha_n^3\right)(1+\delta)\|p_n-x_*\|. \end{aligned} \tag{2.16}$$

Thus, an application of Lemma 1 to (2.15) yields $a_n = \|p_n-u_n\| \to 0$ as $n \to \infty$. Also, since $\|u_n-x_*\| \leq \|p_n-u_n\|+\|p_n-x_*\|$, we have $\|u_n-x_*\| \to 0$ as $n \to \infty$.

Next, we will prove (ii)⇒(i). Assume that $\|u_n - x_*\| \to 0$ as $n \to \infty$. It follows from CR iteration method (1.6), iteration method (1.7) and condition (1.2) that

$$\begin{aligned}
\|u_{n+1} - p_{n+1}\| &= \left\|\left(1-\alpha_n^1\right) v_n + \alpha_n^1 T v_n - \left(1-\alpha_n^1\right) T p_n - \alpha_n^1 T q_n\right\| \\
&\leq \left(1-\alpha_n^1\right)\|v_n - T p_n\| + \alpha_n^1 \|T v_n - T q_n\| \\
&\leq \left(1-\alpha_n^1\right)\left\|\left(1-\alpha_n^2\right) T u_n + \alpha_n^2 T y_n - T p_n\right\| + \alpha_n^1 \delta \|v_n - q_n\| \\
&= \left(1-\alpha_n^1\right)\left\|\left(1-\alpha_n^2\right) T u_n + \alpha_n^2 T y_n - \left(1-\alpha_n^2+\alpha_n^2\right) T p_n\right\| \\
&\quad +\alpha_n^1 \delta \left\|\left(1-\alpha_n^2\right) T u_n + \alpha_n^2 T y_n - \left(1-\alpha_n^2\right) T p_n - \alpha_n^2 T r_n\right\| \\
&\leq \left(1-\alpha_n^1\right)\left(1-\alpha_n^2\right)\|T u_n - T p_n\| + \left(1-\alpha_n^1\right)\alpha_n^2 \|T y_n - T p_n\| \\
&\quad +\alpha_n^1 \delta \left(1-\alpha_n^2\right)\|T u_n - T p_n\| + \alpha_n^1 \delta \alpha_n^2 \|T y_n - T r_n\| \\
&\leq \left(1-\alpha_n^1\right)\left(1-\alpha_n^2\right)\delta\|u_n - p_n\| + \left(1-\alpha_n^1\right)\alpha_n^2 \delta \|y_n - p_n\| \\
&\quad +\alpha_n^1 \delta \left(1-\alpha_n^2\right)\delta\|u_n - p_n\| + \alpha_n^1 \delta \alpha_n^2 \delta \|y_n - r_n\| \\
&= \left\{\left(1-\alpha_n^1\right)\left(1-\alpha_n^2\right)\delta + \alpha_n^1 \delta \left(1-\alpha_n^2\right)\delta\right\}\|u_n - p_n\| \\
&\quad + \left(1-\alpha_n^1\right)\alpha_n^2 \delta \left\|\left(1-\alpha_n^3\right) u_n + \alpha_n^3 T u_n - \left(1-\alpha_n^3+\alpha_n^3\right) p_n\right\| \\
&\quad +\alpha_n^1 \delta \alpha_n^2 \delta \left\|\left(1-\alpha_n^3\right) u_n + \alpha_n^3 T u_n - \left(1-\alpha_n^3\right) p_n - \alpha_n^3 T p_n\right\| \\
&\leq \left\{\left(1-\alpha_n^1\right)\left(1-\alpha_n^2\right)\delta + \alpha_n^1 \delta \left(1-\alpha_n^2\right)\delta\right\}\|u_n - p_n\| \\
&\quad + \left(1-\alpha_n^1\right)\alpha_n^2 \delta \left(1-\alpha_n^3\right)\|u_n - p_n\| + \left(1-\alpha_n^1\right)\alpha_n^2 \delta \alpha_n^3 \|T u_n - p_n\| \\
&\quad +\alpha_n^1 \delta \alpha_n^2 \delta \left(1-\alpha_n^3\right)\|u_n - p_n\| + \alpha_n^1 \delta \alpha_n^2 \delta \alpha_n^3 \|T u_n - T p_n\| \\
&\leq \left\{\left(1-\alpha_n^1\right)\left(1-\alpha_n^2\right)\delta + \alpha_n^1 \delta \left(1-\alpha_n^2\right)\delta\right. \\
&\quad + \left(1-\alpha_n^1\right)\alpha_n^2 \delta \left(1-\alpha_n^3\right) + \left(1-\alpha_n^1\right)\alpha_n^2 \delta \alpha_n^3 \\
&\quad \left.+\alpha_n^1 \delta \alpha_n^2 \delta \left(1-\alpha_n^3\right) + \alpha_n^1 \delta \alpha_n^2 \delta \alpha_n^3 \delta\right\}\|u_n - p_n\| \\
&\quad + \left(1-\alpha_n^1\right)\alpha_n^2 \delta \alpha_n^3 \|u_n - T u_n\| \\
&= \left\{\left(1-\alpha_n^1\right)\left(1-\alpha_n^2\right)\delta + \alpha_n^1 \delta \left(1-\alpha_n^2\right)\delta\right. \\
&\quad + \left(1-\alpha_n^1\right)\alpha_n^2 \delta \left(1-\alpha_n^3\right) + \left(1-\alpha_n^1\right)\alpha_n^2 \delta \alpha_n^3 \\
&\quad \left.+\alpha_n^1 \delta \alpha_n^2 \delta \left[1-\alpha_n^3 (1-\delta)\right]\right\}\|u_n - p_n\| \\
&\quad + \left(1-\alpha_n^1\right)\alpha_n^2 \delta \alpha_n^3 \|u_n - T u_n\|. \qquad (2.17)
\end{aligned}$$

or,

$$\begin{aligned}
\|u_{n+1} - p_{n+1}\| &\leq \left\{\left[1-\alpha_n^1 (1-\delta)\right]\left(1-\alpha_n^2\right)\delta + \left(1-\alpha_n^1\right)\alpha_n^2 \delta\right. \\
&\quad \left.+\alpha_n^1 \delta \alpha_n^2 \delta \left[1-\alpha_n^3 (1-\delta)\right]\right\}\|u_n - p_n\| \\
&\quad + \left(1-\alpha_n^1\right)\alpha_n^2 \delta \alpha_n^3 \|u_n - T u_n\|. \qquad (2.18)
\end{aligned}$$

Since $\delta \in [0,1)$, $\alpha_n^i \in [0,1]$ for all $n \in \mathbb{N}$ and for each $i \in \{1,2,3\}$,

$$1 - \alpha_n^2 (1-\delta) < 1, \qquad (2.19)$$

$$1 - \alpha_n^3 (1-\delta) < 1. \qquad (2.20)$$

By use of inequalities (2.19) and (2.20) in (2.18), we get

$$\begin{aligned}
\|u_{n+1} - p_{n+1}\| &\leq \left[1-\alpha_n^1 (1-\delta)\right]\|u_n - p_n\| \\
&\quad + \left(1-\alpha_n^1\right)\alpha_n^2 \delta \alpha_n^3 \|u_n - T u_n\|. \qquad (2.21)
\end{aligned}$$

Using the fact $x_* = Tx_*$ and triangle inequality for norms, we derive

$$\|u_n - Tu_n\| \leq (1+\delta)\|u_n - x_*\|. \tag{2.22}$$

Hence, (2.21) becomes

$$\begin{aligned} \|u_{n+1} - p_{n+1}\| &\leq \left[1 - \alpha_n^1(1-\delta)\right]\|u_n - p_n\| \\ &\quad + \left(1-\alpha_n^1\right)\alpha_n^2\delta\alpha_n^3(1+\delta)\|u_n - x_*\|. \end{aligned} \tag{2.23}$$

Define

$$\begin{aligned} a_n &= \|u_n - p_n\|, \\ \eta_n &= \alpha_n^1(1-\delta) \in (0,1), \\ \rho_n &= \left(1-\alpha_n^1\right)\alpha_n^2\delta\alpha_n^3(1+\delta)\|u_n - x_*\|. \end{aligned} \tag{2.24}$$

Thus, an application of Lemma 1 to (2.23) yields $a_n = \|u_n - p_n\| \to 0$ as $n \to \infty$. Also, since $\|p_n - x_*\| \leq \|p_n - u_n\| + \|u_n - x_*\|$, we have $\|p_n - x_*\| \to 0$ as $n \to \infty$. □

Theorem 3. *Let S, B and T with fixed point x_* be as in Theorem 1. Let $\left\{\alpha_n^i\right\}_{n=0}^{\infty}$, $i \in \{1,2,3\}$ be real sequences in $[0,1]$ satisfying (i) $\alpha_1 \leq \alpha_n^1 \leq 1$, $\alpha_2 \leq \alpha_n^2 \leq 1$ and $\alpha_3 \leq \alpha_n^3 \leq 1$, for all $n \in \mathbb{N}$ and for some α_1, α_2, $\alpha_3 > 0$. For given $u_0 = p_0 \in S$, consider iterative sequences $\{u_n\}_{n=0}^{\infty}$ and $\{p_n\}_{n=0}^{\infty}$ defined by (1.6) and (1.7), respectively. Then $\{p_n\}_{n=0}^{\infty}$ converges to x_* faster than $\{u_n\}_{n=0}^{\infty}$ does.*

Proof. The following equality was obtained in ([13], Theorem 1)

$$b_n = \|p_0 - x_*\|\delta^{n+1}\left[1 - \alpha_1\left(1 - \delta\left(1 - \alpha_2\alpha_3(1-\delta)\right)\right)\right]^{n+1}. \tag{2.25}$$

Using now (1.7) and (1.2) we have

$$\begin{aligned} \|u_{n+1} - x_*\| &= \left\|\left(1-\alpha_n^1\right)v_n + \alpha_n^1 Tv_n - x_*\right\| \\ &\leq \left(1-\alpha_n^1\right)\|v_n - x_*\| + \alpha_n^1\|Tv_n - x_*\| \\ &\leq \left[\left(1-\alpha_n^1\right) + \alpha_n^1\delta\right]\|v_n - x_*\| \\ &\leq \left[\left(1-\alpha_n^1\right) + \alpha_n^1\delta\right]\left\|\left(1-\alpha_n^2\right)Tu_n + \alpha_n^2 Ty_n - x_*\right\| \\ &\leq \left[\left(1-\alpha_n^1\right) + \alpha_n^1\delta\right]\left(1-\alpha_n^2\right)\|Tu_n - x_*\| \\ &\quad + \left[\left(1-\alpha_n^1\right) + \alpha_n^1\delta\right]\alpha_n^2\|Ty_n - x_*\| \\ &\leq \left[(1-\alpha_n) + \alpha_n^1\delta\right]\left(1-\alpha_n^2\right)\delta\|u_n - x_*\| \\ &\quad + \left[\left(1-\alpha_n^1\right) + \alpha_n^1\delta\right]\alpha_n^2\delta\|y_n - x_*\| \\ &\leq \left[\left(1-\alpha_n^1\right) + \alpha_n^1\delta\right]\left(1-\alpha_n^2\right)\delta\|u_n - x_*\| \\ &\quad + \left[\left(1-\alpha_n^1\right) + \alpha_n^1\delta\right]\alpha_n^2\delta\left(1-\alpha_n^3\right)\|u_n - x_*\| \\ &\quad + \left[\left(1-\alpha_n^1\right) + \alpha_n^1\delta\right]\alpha_n^2\delta\alpha_n^3\delta\|u_n - x_*\| \\ &= \left[\left(1-\alpha_n^1\right) + \alpha_n^1\delta\right]\left\{\left(1-\alpha_n^2\right)\delta + \alpha_n^2\delta\left(1-\alpha_n^3\right) + \alpha_n^2\delta\alpha_n^3\delta\right\}\|u_n - x_*\| \\ &= \left[1 - \alpha_n^1(1-\delta)\right]\left[1 - \alpha_n^2\alpha_n^3(1-\delta)\right]\delta\|u_n - x_*\| \\ &\leq \cdots \\ &\leq \prod_{k=0}^{n}\left[1 - \alpha_k^1(1-\delta)\right]\left[1 - \alpha_k^2\alpha_k^3(1-\delta)\right]\delta\|u_0 - x_*\|. \end{aligned} \tag{2.26}$$

From assumption (i), we obtain

$$\|u_{n+1}-x_*\| \leq \|u_0-x_*\|\,\delta^{n+1}\left[1-\alpha_1\left(1-\delta\right)\right]^{n+1}\left[1-\alpha_2\alpha_3\left(1-\delta\right)\right]^{n+1}. \tag{2.27}$$

Let

$$a_n = \|u_0-x_*\|\,\delta^{n+1}\left[1-\alpha_1\left(1-\delta\right)\right]^{n+1}\left[1-\alpha_2\alpha_3\left(1-\delta\right)\right]^{n+1}. \tag{2.28}$$

Define

$$\begin{aligned}\theta_n &= \frac{a_n}{b_n} = \frac{\|u_0-x_*\|\,\delta^{n+1}\left[1-\alpha_1\left(1-\delta\right)\right]^{n+1}\left[1-\alpha_2\alpha_3\left(1-\delta\right)\right]^{n+1}}{\|p_0-x_*\|\,\delta^{n+1}\left[1-\alpha_1\left(1-\delta\left(1-\alpha_2\alpha_3\left(1-\delta\right)\right)\right)\right]^{n+1}}\\ &= \frac{\left[1-\alpha_1\left(1-\delta\right)\right]^{n+1}\left[1-\alpha_2\alpha_3\left(1-\delta\right)\right]^{n+1}}{\left[1-\alpha_1\left(1-\delta\left(1-\alpha_2\alpha_3\left(1-\delta\right)\right)\right)\right]^{n+1}} \qquad (2.29)\end{aligned}$$

Since $\delta\in(0,1)$ and $\alpha_i\in(0,1)$ for each $i\in\{1,2,3\}$

$$\begin{aligned}\alpha_1 &< 1\\ &\Rightarrow \alpha_1\alpha_2\alpha_3\left(1-\delta\right) < \alpha_2\alpha_3\left(1-\delta\right)\\ &\Rightarrow \alpha_2\alpha_3\left(-1+\delta\right)+\alpha_1\alpha_2\alpha_3\left(1-\delta\right) < 0\\ &\Rightarrow -\alpha_2\alpha_3+\alpha_2\alpha_3\delta+\alpha_1\alpha_2\alpha_3-\alpha_1\alpha_2\alpha_3\delta < 0\\ &\Rightarrow -\alpha_2\alpha_3+\alpha_2\alpha_3\delta+\alpha_1\alpha_2\alpha_3-2\alpha_1\alpha_2\alpha_3\delta < -\alpha_1\alpha_2\alpha_3\delta\\ &\Rightarrow \begin{cases} 1-\alpha_1+\alpha_1\delta-\alpha_2\alpha_3+\alpha_2\alpha_3\delta+\alpha_1\alpha_2\alpha_3-2\alpha_1\alpha_2\alpha_3\delta+\alpha_1\alpha_2\alpha_3\delta^2\\ \qquad < 1-\alpha_1+\alpha_1\delta-\alpha_1\alpha_2\alpha_3\delta+\alpha_1\alpha_2\alpha_3\delta^2\end{cases}\\ &\Rightarrow \begin{cases} 1-\alpha_1+\alpha_1\delta-\alpha_2\alpha_3\left(1-\delta\right)+\alpha_1\alpha_2\alpha_3\left(1-\delta\right)^2\\ \qquad < 1-\alpha_1+\alpha_1\delta-\alpha_1\alpha_2\alpha_3\delta\left(1-\delta\right)\end{cases}\\ &\Rightarrow \begin{cases} 1-\alpha_1\left(1-\delta\right)-\left[1-\alpha_1\left(1-\delta\right)\right]\alpha_2\alpha_3\left(1-\delta\right)\\ \qquad < 1-\alpha_1+\alpha_1\delta\left(1-\alpha_2\alpha_3\left(1-\delta\right)\right)\end{cases}\\ &\Rightarrow \left[1-\alpha_1\left(1-\delta\right)\right]\left[1-\alpha_2\alpha_3\left(1-\delta\right)\right] < 1-\alpha_1\left(1-\delta\left(1-\alpha_2\alpha_3\left(1-\delta\right)\right)\right)\\ &\Rightarrow \frac{\left[1-\alpha_1\left(1-\delta\right)\right]\left[1-\alpha_2\alpha_3\left(1-\delta\right)\right]}{1-\alpha_1\left(1-\delta\left(1-\alpha_2\alpha_3\left(1-\delta\right)\right)\right)} < 1, \qquad (2.30)\end{aligned}$$

and thus, we have

$$\begin{aligned}\lim_{n\to\infty}\frac{\theta_{n+1}}{\theta_n} &= \lim_{n\to\infty}\frac{\frac{\left[1-\alpha_1(1-\delta)\right]^{n+2}\left[1-\alpha_2\alpha_3(1-\delta)\right]^{n+2}}{\left[1-\alpha_1(1-\delta(1-\alpha_2\alpha_3(1-\delta)))\right]^{n+2}}}{\frac{\left[1-\alpha_1(1-\delta)\right]^{n+1}\left[1-\alpha_2\alpha_3(1-\delta)\right]^{n+1}}{\left[1-\alpha_1(1-\delta(1-\alpha_2\alpha_3(1-\delta)))\right]^{n+1}}}\\ &= \lim_{n\to\infty}\frac{\left[1-\alpha_1\left(1-\delta\right)\right]\left[1-\alpha_2\alpha_3\left(1-\delta\right)\right]}{1-\alpha_1\left(1-\delta\left(1-\alpha_2\alpha_3\left(1-\delta\right)\right)\right)}\\ &= \frac{\left[1-\alpha_1\left(1-\delta\right)\right]\left[1-\alpha_2\alpha_3\left(1-\delta\right)\right]}{1-\alpha_1\left(1-\delta\left(1-\alpha_2\alpha_3\left(1-\delta\right)\right)\right)} < 1. \qquad (2.31)\end{aligned}$$

It thus follows from ratio test that $\sum_{n=0}^{\infty}\theta_n<\infty$. Hence, we have $\lim_{n\to\infty}\theta_n=0$ which implies that $\{p_n\}_{n=0}^{\infty}$ is faster than $\{u_n\}_{n=0}^{\infty}$. □

We are now able to establish the following data dependence result.

Theorem 4. *Let $\widetilde{T}$ be an approximate operator of T satisfying condition (1.2). Let $\{p_n\}_{n=0}^{\infty}$ be an iterative sequence generated by (1.7) for T and define an iterative*

sequence $\{\widetilde{p}_n\}_{n=0}^{\infty}$ *as follows*

$$\left\{\begin{array}{l} \widetilde{p}_0 \in S, \\ \widetilde{p}_{n+1} = \left(1-\alpha_n^1\right)\widetilde{T}\widetilde{p}_n + \alpha_n^1\widetilde{T}\widetilde{q}_n, \\ \widetilde{q}_n = \left(1-\alpha_n^2\right)\widetilde{T}\widetilde{p}_n + \alpha_n^2\widetilde{T}\widetilde{r}_n, \\ \widetilde{r}_n = \left(1-\alpha_n^3\right)\widetilde{p}_n + \alpha_n^3\widetilde{T}\widetilde{p}_n,\ n\in\mathbb{N}, \end{array}\right. \tag{2.32}$$

where $\left\{\alpha_n^i\right\}_{n=0}^{\infty}$, $i\in\{1,2,3\}$ *be real sequences in* $[0,1]$ *satisfying (i)* $\frac{1}{2}\leq\alpha_n^1$ *for all* $n\in\mathbb{N}$, *and (ii)* $\sum\limits_{n=0}^{\infty}\alpha_n^1=\infty$. *If* $Tp=p$ *and* $\widetilde{T}\widetilde{p}=\widetilde{p}$ *such that* $\widetilde{p}_n\to\widetilde{p}$ *as* $n\to\infty$, *then we have*

$$\|p-\widetilde{p}\|\leq\frac{5\varepsilon}{1-\delta}, \tag{2.33}$$

where $\varepsilon>0$ *is a fixed number.*

Proof. It follows from (1.2), (1.7), and (2.32) that

$$\begin{aligned} \|r_n-\widetilde{r}_n\| &= \left\|\left(1-\alpha_n^3\right)p_n+\alpha_n^3Tp_n-\left(1-\alpha_n^3\right)\widetilde{p}_n-\alpha_n^3\widetilde{T}\widetilde{p}_n\right\| \\ &\leq \left(1-\alpha_n^3\right)\|p_n-\widetilde{p}_n\|+\alpha_n^3\left\|Tp_n-\widetilde{T}\widetilde{p}_n\right\| \\ &\leq \left(1-\alpha_n^3\right)\|p_n-\widetilde{p}_n\|+\alpha_n^3\left\{\|Tp_n-T\widetilde{p}_n\|+\left\|T\widetilde{p}_n-\widetilde{T}\widetilde{p}_n\right\|\right\} \\ &\leq \left[1-\alpha_n^3(1-\delta)\right]\|p_n-\widetilde{p}_n\|+\alpha_n^3\varepsilon, \end{aligned} \tag{2.34}$$

$$\begin{aligned} \|q_n-\widetilde{q}_n\| &= \left\|\left(1-\alpha_n^2\right)Tp_n+\alpha_n^2Tr_n-\left(1-\alpha_n^2\right)\widetilde{T}\widetilde{p}_n-\alpha_n^2\widetilde{T}\widetilde{r}_n\right\| \\ &\leq \left(1-\alpha_n^2\right)\left\|Tp_n-\widetilde{T}\widetilde{p}_n\right\|+\alpha_n^2\left\|Tr_n-\widetilde{T}\widetilde{r}_n\right\| \\ &\leq \left(1-\alpha_n^2\right)\left\{\|Tp_n-T\widetilde{p}_n\|+\left\|T\widetilde{p}_n-\widetilde{T}\widetilde{p}_n\right\|\right\} \\ &\quad +\alpha_n^2\left\{\|Tr_n-T\widetilde{r}_n\|+\left\|T\widetilde{r}_n-\widetilde{T}\widetilde{r}_n\right\|\right\} \\ &\leq \left(1-\alpha_n^2\right)\delta\|p_n-\widetilde{p}_n\|+\alpha_n^2\delta\|r_n-\widetilde{r}_n\|+\left(1-\alpha_n^2\right)\varepsilon+\alpha_n^2\varepsilon, \end{aligned} \tag{2.35}$$

$$\begin{aligned} \|p_{n+1}-\widetilde{p}_{n+1}\| &= \left\|\left(1-\alpha_n^1\right)Tp_n+\alpha_n^1Tq_n-\left(1-\alpha_n^1\right)\widetilde{T}\widetilde{p}_n-\alpha_n^1\widetilde{T}q_n\right\| \\ &\leq \left(1-\alpha_n^1\right)\left\|Tp_n-\widetilde{T}\widetilde{p}_n\right\|+\alpha_n^1\left\|Tq_n-\widetilde{T}\widetilde{q}_n\right\| \\ &\leq \left(1-\alpha_n^1\right)\left\{\|Tp_n-T\widetilde{p}_n\|+\left\|T\widetilde{p}_n-\widetilde{T}\widetilde{p}_n\right\|\right\} \\ &\quad +\alpha_n^1\left\{\|Tq_n-T\widetilde{q}_n\|+\left\|T\widetilde{q}_n-\widetilde{T}\widetilde{q}_n\right\|\right\} \\ &\leq \left(1-\alpha_n^1\right)\left\{\delta\|p_n-\widetilde{p}_n\|+\varepsilon\right\}+\alpha_n^1\left\{\delta\|q_n-\widetilde{q}_n\|+\varepsilon\right\} \\ &= \left(1-\alpha_n^1\right)\delta\|p_n-\widetilde{p}_n\|+\alpha_n^1\delta\|q_n-\widetilde{q}_n\|+\left(1-\alpha_n^1\right)\varepsilon+\alpha_n^1\varepsilon. \end{aligned} \tag{2.36}$$

Combining (2.34), (2.35), and (2.36)

$$\begin{aligned} \|p_{n+1}-\widetilde{p}_{n+1}\| &\leq \left\{\left(1-\alpha_n^1\right)\delta+\alpha_n^1\delta\left\{\left(1-\alpha_n^2\right)\delta+\alpha_n^2\delta\left[1-\alpha_n^3(1-\delta)\right]\right\}\right\}\|p_n-\widetilde{p}_n\| \\ &\quad +\alpha_n^1\delta\alpha_n^2\delta\alpha_n^3\varepsilon+\alpha_n^1\delta\left(1-\alpha_n^2\right)\varepsilon+\alpha_n^1\delta\alpha_n^2\varepsilon+\left(1-\alpha_n^1\right)\varepsilon+\alpha_n^1\varepsilon \end{aligned} \tag{2.37}$$

Since $\delta\in(0,1)$, $\alpha_n^i\in[0,1]$ for each $i\in\{1,2,3\}$ and for all $n\in\mathbb{N}$,

$$1-\alpha_n^3(1-\delta)<1, \tag{2.38}$$

$$1-\alpha_n^2\left(1-\delta\right)<1, \tag{2.39}$$

$$\alpha_n^2\alpha_n^3\delta^2<1, \tag{2.40}$$

$$\left(1-\alpha_n^2\right)\delta<1, \tag{2.41}$$

$$\alpha_n^2\delta<1, \tag{2.42}$$

and by assumption (i) we have

$$1-\alpha_n^1\leq\alpha_n^1. \tag{2.43}$$

Thus, an application of inequalities (2.38), (2.39), (2.40), (2.41), (2.42) and (2.43) to (2.37) yields

$$\left\|p_{n+1}-\widetilde{p}_{n+1}\right\|\leq\left[1-\alpha_n^1\left(1-\delta\right)\right]\left\|p_n-\widetilde{p}_n\right\|+\alpha_n^1\left(1-\delta\right)\frac{5\varepsilon}{1-\delta}. \tag{2.44}$$

Let us denote

$$a_n:=\left\|p_n-\widetilde{p}_n\right\|,\ \mu_n:=\alpha_n^1\left(1-\delta\right)\in\left(0,1\right). \tag{2.45}$$

It follows from Lemma 2 that

$$0\leq\limsup_{n\to\infty}\left\|p_n-\widetilde{p}_n\right\|\leq\limsup_{n\to\infty}\frac{5\varepsilon}{1-\delta}. \tag{2.46}$$

From Theorem 1 we know that $\lim_{n\to\infty}p_n=x_*$. Thus, using this fact together with the assumption $\lim_{n\to\infty}\widetilde{p}_n=u_*$ we obtain

$$\left\|x_*-u_*\right\|\leq\frac{5\varepsilon}{1-\delta}. \tag{2.47}$$

□

References

[1] Agarwal, R., O Regan, D. & Sahu, D. 2007. Iterative construction of fixed points of nearly asymptotically nonexpansive mappings. Journal of Nonlinear and Convex Analysis 8: 61-79.

[2] Babu, G. V. R. & Prasad, K. N. V. V. V. 2006. Mann iteration converges faster than Ishikawa iteration for the class of Zamfirescu operators. Fixed Point Theory and Applications 2006(1).

[3] Berinde, V. 2007. Iterative Approximation of Fixed Points. Springer, Berlin.

[4] Berinde, V. 2004. Picard iteration converges faster than the Mann iteration in the class of quasi-contractive operators. Fixed Point Theory Appl. 2: 97-105.

[5] Berinde, V. 2004. Comparing Krasnoselskij and Mann iterations for Lipschitzian generalized pseudocontractive operators. Proceed. of Int. Conf. On Fixed Point Theory, Univ. of Valencia. Yokohama Publishers.

[6] Berinde, V. & Berinde, M. 2005. The fastest Krasnoselskij iteration for approximating fixed points of strictly pseudo-contractive mappings. Carpathian J. Math 21(1-2): 13-20.

[7] Chang, S. S., Cho, Y. J. & Kim, J. K. 2003. The equivalence between the convergence of modified Picard, modified Mann, and modified Ishikawa iterations. Mathematical and Computer Modelling 37(9): 985-991.

[8] Chugh, R. & Kumar, V. 2012. Stability of hybrid fixed point iterative algorithms of Kirk-Noor type in normed linear space for self and nonself operators. International Journal of Contemporary Mathematical sciences 7: 1165-1184.

[9] Chugh, R., Kumar, V. & Kumar, S. 2012. Strong Convergence of a new three step iterative scheme in Banach spaces. American Journal of Computational Mathematics 2: 345-357.

[10] Hussain, N., Chugh, R., Kumar, V. & Rafiq, A. 2012. On the rate of convergence of Kirk-type iterative schemes. Journal of Applied Mathematics 2012.

[11] Hussain, N., Rafiq, A., Damjanović, B. & Lazović, R. 2011. On rate of convergence of various iterative schemes. Fixed Point Theory and Applications 2011(1): 1-6.

[12] Ishikawa, S. 1974. Fixed points by a new iteration method. Proceedings of the American Mathematical Society 44: 147-150.
[13] Karahan, I. & Ozdemir, M. 2013. A general iterative method for approximation of fixed points and their applications. Advances in Fixed Point Theory 3(3).
[14] Karakaya, V., Doğan, K., Gürsoy, F. & Ertürk, M. 2013. Fixed Point of a New Three-Step Iteration Algorithm under Contractive-Like Operators over Normed Spaces. Abstract and Applied Analysis 2013.
[15] Kirk, W. 1971. On successive approximations for nonexpansive mappings in Banach spaces. Glasgow Math. J. 12:6-9.
[16] Krasnosel'skii, M.A. 1955. Two remarks on the method of successive approximations. Uspekhi Matematicheskikh Nauk 10: 123-127.
[17] Mann, W.R. 1953. Mean value methods in iteration. Proceedings of the American Mathematical Society 4: 506-510.
[18] Noor, M.A. 2000. New approximation schemes for general variational inequalities. Journal of Mathematical Analysis and Applications 251: 217-229.
[19] Olatinwo, M. 2009. Some stability results for two hybrid fixed point iterative algorithms in normed linear space. Matematique Vesnik 61: 247-256.
[20] Olatinwo, M., Owojori, O. & Imoru, C. 2006. Some stability results for fixed point iteration processes. Aus. J. Math. Anal. Appl. 3: 1-7.
[21] Phuengrattana, W. & Suantai, S. 2012. Comparison of the Rate of Convergence of Various Iterative Methods for the Class of Weak Contractions in Banach Spaces. Thai Journal of Mathematics 11(1): 217-226.
[22] Phuengrattana, W. & Suantai, S. 2011. On the rate of convergence of Mann, Ishikawa, Noor and SP-iterations for continuous functions on an arbitrary interval. Journal of Computational and Applied Mathematics 235: 3006-3014.
[23] Picard, E., (1890). "Mémoire sur la théorie des équations aux dérivées partielles et la méthode des approximations successives", Journal de Mathématiques pures et appliquées, 6:145-210.
[24] Popescu, O. 2007. Picard iteration converges faster than Mann iteration for a class of quasicontractive operators. Math. Commun. 12: 195-202.
[25] Rhoades, B. E. & Şoltuz, S. M. 2004. The equivalence between the convergences of Ishikawa and Mann iterations for an asymptotically nonexpansive in the intermediate sense and strongly successively pseudocontractive maps. Journal of mathematical analysis and applications 289(1): 266-278.
[26] Rhoades, B. & Soltuz, S.M. 2004. The equivalence between Mann–Ishikawa iterations and multistep iteration. Nonlinear Analysis: Theory, Methods & Applications 58: 219-228.
[27] Rhoades, B. E. 1976. Comments on two fixed point iteration methods. J. Math. Anal. Appl. 56(2): 741-750.
[28] Sahu, D. R. 2011. Applications of the S-iteration process to constrained minimization problems and split feasibility problems. Fixed Point Theory 12(1): 187-204.
[29] Schaefer, H. 1957. Über die Methode sukzessiver Approximationen. Jahresbericht der Deutschen Mathematiker-Vereinigung 59: 131-140.
[30] Soltuz, S.M. & Grosan, T. 2008. Data dependence for Ishikawa iteration when dealing with contractive like operators. Fixed Point Theory and Applications 2008: 242916(1-7).
[31] Sun, H. & Xue, Z. 2013. Reconsiderations on the Equivalence of Convergence between Mann and Ishikawa Iterations for Asymptotically Pseudocontractive Mappings. Journal of Applied Mathematics 2013.
[32] Thianwan, S. 2009. Common fixed points of new iterations for two asymptotically nonexpansive nonself-mappings in a Banach space. Journal of Computational and Applied Mathematics 224: 688-695.
[33] Weng, X. 1991. Fixed point iteration for local strictly pseudocontractive mapping. Proc. Amer. Math. Soc. 113: 727-731.
[34] Xue, Z. 2008. The comparison of the convergence speed between Picard, Mann, Krasnoselskij and Ishikawa iterations in Banach spaces. Fixed Point Theory Appl. 2008.
[35] Yuan, Q. & Rhoades, B.E. 2008. Comments on the rate of convergence between Mann and Ishikawa iterations applied to Zamfirescu operators. Fixed Point Theory and Applications 2008.

Department of Mathematical Engineering, Yildiz Technical University, Davutpasa Campus, Esenler, 34210 İstanbul, Turkey
E-mail address: vkkaya@yildiz.edu.tr; vkkaya@yahoo.com
URL: http://www.yarbis.yildiz.edu.tr/vkkaya

Department of Mathematics, Yildiz Technical University, Davutpasa Campus, Esenler, 34220 Istanbul, Turkey
E-mail address: faikgursoy02@hotmail.com; fgursoy@yildiz.edu.tr
URL: http://www.yarbis.yildiz.edu.tr/fgursoy

Department of Mathematics, Yildiz Technical University, Davutpasa Campus, Esenler, 34220 Istanbul, Turkey
E-mail address: merturk3263@gmail.com
URL: http://www.yarbis.yildiz.edu.tr/merturk